

## OS04lite ACL Reconstruction

### Further information

You can get more information about this procedure from [www.aboutmyhealth.org](http://www.aboutmyhealth.org)

You can get information locally from the Senior Nurse at the Horder Centre,  
St John's Road, Crowborough,  
East Sussex, TN6 1XP.  
Tel: 01892 665577  
[info@hordercentre.co.uk](mailto:info@hordercentre.co.uk)

Tell us how useful you found this document at [www.patientfeedback.org](http://www.patientfeedback.org)



[www.rcsed.ac.uk](http://www.rcsed.ac.uk)



This document will give you information about an ACL reconstruction. If you have any questions, you should ask your GP or other relevant health professional.

### What is the anterior cruciate ligament?

The anterior cruciate ligament (ACL) is one of the important ligaments that stabilise your knee joint (see figure 1). If you have torn (ruptured) this ligament, the knee can collapse or 'give way' when making twisting or turning movements.




Figure 1  
Internal structure of the right knee

### How does an ACL rupture happen?

An ACL rupture happens as a result of a twisting injury to the knee. The common causes are football and skiing injuries. You can injure other parts of your knee at the same time, such as tearing a cartilage or damaging the joint surface.

### What are the benefits of surgery?

If your ACL reconstruction is successful, your knee should not give way any more. This will allow you to be more active and return to some or all of your sporting activities.

### Are there any alternatives to surgery?

Your physiotherapist can give you exercises to strengthen and improve the co-ordination of the muscles in your thigh. Wearing a knee brace can sometimes help during sports activities.

### What does the operation involve?

A variety of anaesthetic techniques are possible. The operation usually takes between an hour and an hour and a half. Your surgeon will make one or more cuts around your knee. Some surgeons perform the operation by arthroscopy ('keyhole' surgery) using a camera to see inside the knee.

Your surgeon will replace the ACL with a piece of suitable tissue from elsewhere in the body. The ends of the replacement ligament are fixed with special screws or anchors into 'tunnels' drilled in the bone.

### What complications can happen?

#### 1 General complications

- Pain
- Bleeding
- Infection of the surgical site (wound)
- Unsightly scarring
- Blood clots
- Difficulty passing urine

#### 2 Specific complications

- Break of the kneecap
- Damage to nerves around the knee
- Infection in the knee joint
- Discomfort in the front of the knee
- Loss of knee movement
- The knee keep giving way
- Severe pain, stiffness and loss of use of the knee (Complex Regional Pain Syndrome)

### How soon will I recover?

You should be able to go home the same day or the day after.

Your surgeon may want you to wear a knee brace for a few weeks after the operation. Once your knee is settling down you will need to start intensive physiotherapy treatment which may continue for as long as six months.

Regular exercise should help you to return to normal activities as soon as possible. Before you start exercising, you should ask a member of the healthcare team or your GP for advice.

It is unlikely that your knee will ever be quite as good as it was before the original injury.

## Summary

If your knee continually gives way after an ACL rupture, reconstruction offers the chance of improving the stability of your knee in everyday life and in sporting activities. You may be able to return to a level of sport that otherwise would not be possible.

## Acknowledgements

Author: Mr Stephen Milner DM FRCS (Tr. & Orth.)  
Illustrations: Medical Illustration Copyright © 2011  
Nucleus Medical Art. All rights reserved.  
www.nucleusinc.com

**This document is intended for information purposes only and should not replace advice that your relevant health professional would give you.**